MINNEAPOLIS PLASTIC SURGERY, LTD.

763-545-0443 (24 hours)

If answering service does not pick up call 952-653-0784. In an EMERGENCY call 911

AFTER SURGERY INSTRUCTIONS BREAST SURGERY

- Take your medications as prescribed. See separate page.
- Call your doctor if you notice <u>progressive</u> swelling (some swelling is to be expected), bleeding, or soreness, particularly if you notice the difference only on one breast.
- It is necessary that you stay quiet for the next 24 hours, as excessive activity may cause bleeding to occur. Walking is OK; long walks or treadmills are NOT! Flex your calves to keep blood moving.
- Sleep with your upper body elevated for 3 weeks. This will help decrease swelling and soreness. A sofa cushion under the mattress works better than several pillows. A comfortable recliner is OK too.
- You may drink liquids when you arrive home. Later, if you are not nauseated, you may have a light meal.
- You may shower tomorrow. Do not go in any hot tubs, tub baths, pools, or Jacuzzis until OK'd by the doctor (minimum of 3 weeks). Do NOT remove Band-Aids or tapes before showering.
- ❖ You may apply dry cloth-protected ice packs to breasts as needed. **DO NOT** use ice on the lower breast areas if you had breast lift surgery. **DO NOT** apply ice directly to skin or blisters/frostbite can occur!
- ❖ <u>DO NOT smoke</u> (including second-hand smoke) or use nicotine products of any kind for 2 weeks before and after breast surgery. This is even more critical if you had breast lift surgery.
- ❖ Keep the dressings on your breasts intact until you see your doctor. **DO NOT** remove Band-Aids or tapes.
- ❖ **DO NOT** lift any objects greater than five (5) lbs. for 2 weeks.

I acknowledge receipt of these after-surgery instructions.

- ❖ **DO NOT** engage in any strenuous activity that involves the use of your arms, i.e. vacuuming, heavy lifting, or raising your arms above your head for 2-3 weeks.
- ❖ **DO NOT** drink any alcoholic beverages (<u>beer and wine included</u>) for 5 days or while taking pain medication. Alcohol, medications and/or anesthetic drugs may interfere with good judgment.
- ❖ **DO NOT** drive a car or operate machinery for the next 48 hours or while using pain pills.
- Leave Exparel bracelet on for 4 days, as this is how long this medication is in your body.
- Someone from Minneapolis Plastic Surgery will call you this evening to inquire about your condition and answer any questions you may have.
- Do not hesitate to call the doctor's office if you have any questions about your recovery. PHONE: 763-545-0443

(signature)	(date and time)